

Advance Publication by J-STAGE

Japanese Journal of Infectious Diseases

Evaluation of the anti-adenoviral activity of ozonated alcohol disinfectant - ALTANT

Nozomu Hanaoka, Naomi Nojiri, Kenichiro Takahashi, Eiichi Yoshida, and Tsuguto
Fujimoto

Received: February 17, 2020. Accepted: March 31, 2020.

Published online: April 30, 2020.

DOI: 10.7883/yoken.JJID.2020.058

Advance Publication articles have been accepted by JJID but have not been copyedited
or formatted for publication.

Title page

Title

Evaluation of the anti-adenoviral activity of ozonated alcohol disinfectant - ALTANT

Author

Nozomu Hanaoka^{1*}, Naomi Nojiri¹, Kenichiro Takahashi¹, Eiichi Yoshida², Tsuguto

Fujimoto¹

Affiliation

¹ Laboratory Diagnosis Division, Infectious Disease Surveillance Center, National Institute of Infectious Diseases, 1-23-1 Toyama, Shinjyuku-ku, Tokyo 162-8640, Japan

² E-TECH Co., Ltd., Kobe-Incubation-Office 316, 9-1 Minatojima, Chuo-ku, Kobe, Hyogo 654-0151, Japan

***Corresponding author**

Nozomu Hanaoka, Ph.D.

E-Mail: nozomu@nih.go.jp

Phone: 81-3-5285-1111

Fax: 81-3-5285-1129

Address: Laboratory Diagnosis Division, Infectious Disease Surveillance Center,
National Institute of Infectious Diseases, 1-23-1 Toyama, Shinjyu-ku, Tokyo 162-8640,
Japan

Key words

Human masrtadenovirus, disinfectant, ozonated alcohol

Running head

Anti-adenoviral activity of ALTANT

花岡希^{1*}、野尻直未¹、高橋健一郎¹、吉田英一²、藤本嗣人¹

¹ 国立感染症研究所 感染症疫学センター 第四室

² 株式会社 E テック。

*責任著者

花岡希

〒162-8640 東京都新宿区戸山 1-23-1

電話: 81-3-5285-1111

Fax: 81-3-5285-1129 .

E-Mail: nozomu@nih.go.jp

Key word

Human mastadenovirus, 消毒薬, オゾン化アルコール.

Running head

アルタントの抗アデノウイルス効果

Summary

Seven species (A–G) of human mastadenovirus (HAdV) are known, and more than 100 types have been reported. HAdV is highly resistant to common hand sanitizers. Epidemic keratoconjunctivitis and pharyngoconjunctival fever caused by HAdV can be explosively transmitted in a confined space, resulting in outbreaks such as nosocomial infections. Given the absence of an antiviral agent for HAdV infection, it is important to prevent the spread of infection using disinfectants. The bactericidal and virucidal activities of ozone have already been known widely. ALTANT is an ozonated alcohol preparation developed by E-TECH Co., Ltd. (Kobe, Hyogo, Japan).

In this study, ALTANT was mixed with different types of HAdV at a ratio of 9:1, and the viability of the HAdV was determined after instantaneous reactions for varying time periods (flash (< 3 seconds) to 5 minutes) using TCID₅₀ assay. The assay results demonstrated that the viability of the HAdV decreased by 1/10 to 1/100 within 1 minute after the reaction, albeit slight differences in the reactivity observed among HAdV types. The viability of the HAdV decreased by a factor of $>4\log_{10}$, and the virus was completely killed in 3 minutes. This study demonstrated that ALTANT has a potent HAdV disinfection effect.

Main Text

Introduction

Human mastadenovirus (HAdV) is a nonenveloped DNA virus and is classified into seven species, i.e., A–G. Currently, more than 100 different types have been reported for HAdV (1). HAdV has been reported to be highly resistant to disinfectants (2,3), particularly to alcoholic preparations (4,5). Chlorine is effective for the disinfection of HAdV (2). Pharyngoconjunctival fever (PCF), also known as swimming pool fever, and epidemic keratoconjunctivitis (EKC) are typical acute and highly infectious adenoviral diseases. Relevant laws prescribe that swimming pool water must be disinfected with free residual chlorine at a concentration between 0.4 mg/L and 1.0 mg/L. Sodium hypochlorite is a recommendable chlorine-based disinfectant; however, it is difficult to use for the disinfection of hands, eyes, and metallic instruments owing to the mucosal irritation and metal corrosion properties of chlorine gas.

HAdV-B11 associated with hemorrhagic cystitis is a major concern in hematopoietic stem cell transplantation because it causes secondary infection due to the infection of the virus in the environment (6–9). Post transplantation infection caused by adenovirus species B or C, is a problem worldwide (10–16). Adenoviruses associated with these infections are known to be explosively transmitted in confined spaces such as hospitals.

Patients with HAdV have been reported to discharge infectious viruses in their saliva and urine for an extended period of time (17). Adenoviruses have high infectivity, and HAdV can propagate even when only one virion is present in a culture medium (17,18). Considering that no anti-HAdV drugs are presently in clinical use, disinfectants are important for preventing outbreak of HAdV (1,10) and it is important to evaluate the virucidal effects of disinfectants on HAdVs.

The bactericidal and virucidal activities of ozone have already been known widely (19). Disinfection with ozonated water solutions has been used for water sterilization in water supply and sewerage systems (19). Ozone has been demonstrated to be effective for enteric adenovirus types 40 and 41 in sewage treatment (20,21). In water, ozone is relatively stable in acidic solutions, but it rapidly decomposes at an elevated temperature or pH. Under such decomposing conditions, ozone is hydrolyzed to generate hydroxyl radical ($\bullet\text{OH}$), hydroperoxyl radical (HO_2), or hydrogen peroxide (H_2O_2). In particular, hydroxy radicals have a higher oxidizing capacity than ozone molecules, and they contribute greatly to the inactivation of bacteria and viruses (19). Besides high antibacterial and antiviral effects, the advantages of using ozone include no generation of organochlorine compounds, such as trihalomethane; no persistence; easy removal of iron and manganese; and deodorization effect. Its disadvantages include transient efficacy due

to no persistence, high cytotoxicity even at relatively low concentrations, and high corrosiveness, particularly to rubber and plastics.

ALTANT is an ozonated alcohol preparation developed by E-TECH Co., Ltd. (Kobe, Hyogo, Japan) (<http://www.e-teck.co.jp/>). ALTANT is a disinfectant in which ozone is stably retained in alcohol. ALTANT is at least as effective as other disinfectants, such as peracetic acid, glutaral, and phytoral (<http://www.e-teck.co.jp/>). Its safety has been evaluated in various tests, including acute oral toxicity, eye irritancy, skin irritancy, mutagenicity, and oral mucosal irritation tests (Japan Food Research Laboratories; No.17094693001-0101). ALTANT has been patented in several countries, including Japan, the United States, the EU, South Korea, and Taiwan. It has been demonstrated (Kitakanhatu 2017_0026 and Hokuseihatu 2017_0161) to have antiseptic effects against *Bacillus subtilis* (ATCC 6633) and *Feline calicivirus* (strain F-9; ATCC VR-782), which is a norovirus surrogate. ALTANT is expected to exhibit a broad antiviral effect.

Several disinfectants have been tested for their efficacy against adenoviruses; however, only a few types includes type 3 of species B and type 5 of species C have been used in these tests (22,23). Recently, recombinant adenoviruses have been prevalent worldwide (1). However, many recombinant adenoviruses have not been fully characterized, and their susceptibility to disinfectants remains unknown. Although there

are several types of HAdVs worldwide, the prevalent types differ among countries and regions (Infectious Agents Surveillance Report (IASR): <https://www.niid.go.jp/niid/en/865-iasr/7390-449te.html>). Moreover, HAdVs of different serotypes have been reported to show different responses to a disinfectant (2,24). Disinfectants should be evaluated for their effects on specific HAdV types prevalent in Japan.

The present study evaluated the anti-adenoviral effects of ALTANT against 14 types of HAdVs that cause respiratory system, ophthalmic, genito-urinary tract, and systemic infections.

Materials and Methods

Virus preparation

A549 cells (CCL-185, American Type Culture Collection (ATCC)) was used as host cells for the adenovirus culture. A549 cells were grown with Eagle's Minimum Essential Medium with Earle's salts (Eagle's MEM, Sigma-Aldrich Japan, Tokyo, Japan) comprising 10 % fetal bovine serum (FBS; Biowest, Nuaille, France), 1% L-alanine/L-glutamine (200 mmol/L) (Wako Pure Chemical Industries (Wako, Osaka, Japan), 0.2%

gentamicin sulfate solution (50 mg/ml) (Wako), and 0.1% amphotericin B (Wako).

HAdVs were inoculated in A549 cells with the same medium (except with 5% FBS instead of 10% FBS) and incubated until cytopathic effects (CPE) were visually observed with a WRAYCAM (WRAYMER Inc., Osaka, Japan) camera using an inverted microscope CKX53 (Olympus Inc., Tokyo, Japan). Reference adenovirus types were either purchased from ATCC or reference strains as reported previously (25). Information regarding endemic the types of HAdV in Japan is obtained from IASR. In the present study, HAdV types were selected based on IASR data. A total of 14 adenovirus types were used as follows: HAdV-1, HAdV-2, HAdV-3, HAdV-4, HAdV-5, and HAdV-6 responsible for PCF; HAdV-7 responsible for ARDS; HAdV-11 responsible for HC; HAdV-37, HAdV-53, HAdV-54, HAdV-56, HAdV-64, and HAdV-85 responsible for EKC. HAdV-85 has been recently reported as a causative agent of conjunctivitis in Japan (25). For preparing the virus stock solution, intact adenovirus was roughly purified from the cell culture, and complete CPE was observed as follows. The virus culture supernatant was centrifuged at 1,500×g for 2 minutes, and the supernatant was collected and dispensed into several tubes and then preserved at −80°C until further evaluation.

Virus quantification and TCID50 assay

Virus quantification was performed using real-time polymerase chain reaction (PCR) as described previously (26). HAdV titration, which was adjusted to 1×10^5 copies, was performed using microtiter plates by two-fold serial dilutions. The diluted virus sample was inoculated into each of the 96 wells comprising confluent A549 cells at 34°C with 5% CO₂, and the appearance of CPE was observed daily for 20 days. The Spearman-Kärber method was used to calculate the median tissue culture infective dose (TCID₅₀)/ml [(27,28)].

Effects of ALTANT on the A549 cell line

ALTANT was provided by E-TECH Co., Ltd.. To determine the effects of ALTANT on A549 cells, dilution series of ALTANT was performed using a cell growth medium of A549 cells. ALTANT was mixed with the medium at varying ratios as follows: 1:2, 1:5, 1:10, 1:50, 1:100, 1:1000, and 1:5000, respectively. The diluted solution was applied to confluent A549 cells in a 96-well plate and was incubated at 34°C for 7 days. Cell growth was recorded every day, and cells passage was performed at day 7 to evaluate the effects on cell growth. Floating A549 cells treated with trypsin were also evaluated using the same method. The effects of isopropyl alcohol, which is the base material of ALTANT, were also evaluated as described above.

Neutralizer of ALTANT

To neutralize ALTANT, mixtures of ALTANT and the cell growth medium with 0.1%, 0.5% and 1% sodium thiosulphate were evaluated. Each solution was applied to confluent A549 cells and incubated at 34°C for 7 days. Cell growth was analyzed as described above.

Evaluation of anti-adenoviral activity of ALTANT

Based on the European Committee for Standardization (CEN: <https://www.cen.eu/Pages/default.aspx>) EN 14476 (Chemical disinfectants and antiseptics) and previously described methods (24,29), a modified assay was performed as follows.

The evaluation assay was performed using the “endpoint” method. To quantify the amount of virus, A549 cells were propagated in 96-well cell culture plates. To investigate whether ALTANT has stabilizing or disinfection activity for HAdV, the mixture of HAdV was neutralized with ALTANT, diluted to 1^{-0} – 10^{-20} by two-fold serial dilution, inoculated into A549 cells in 96-well plates, and then the A549 cells were incubated at 34°C for 21 days. The time (day) and rate of CPE appearance were determined (18,24,29). One row, including 8 wells of the 96 well plate, was used for one

dilution. The detection of CPE-positive rows was determined when CPE was observed in $\geq 5/8$ wells. To calculate the TCID₅₀ of the endpoint of CPE observation, all CPE-positive line was regarded as 8/8 among all the tests. This assay was performed as follows: 900 μL of ALTANT and 100 μL of the virus stock solution (1×10^5 copies) was mixed by vortexing for 2 seconds; then this mixture was kept standing at room temperature for the following time periods— flash (within 3 seconds), 10 seconds, 30 seconds, 1 minute, 3 minutes, and 5 minutes; 10 μL of the reacted solution was neutralized by 990 μL of 0.5 % sodium thiosulphate. Subsequently, 100 μL of the neutralized mixture was diluted to 2^{-1} – 2^{-20} by two-fold serial dilution with 100 μL fresh medium. Then 100 μL of diluted solution was added to confluent A549 cells filled with 100 μL fresh medium at each well in the 96-well plate; this plate was incubated at 34°C for 21 days, and CPE was observed regularly. All the assays were performed twice independently. Eagle's MEM medium was used instead of ALTANT as the untreated control.

Results

Cytotoxicity and neutralization of ALTANT

ALTANT did not affect cell growth at a dilution of 1/1000 in the MEM cell growth

medium but inhibited cell growth at a dilution of 1/500. In the neutralization test of ALTANT using sodium thiosulfate, the cell toxicity of ALTANT on the growth of A549 cells was abolished when it was mixed at a ratio of 1:100 with the MEM cell growth medium comprising 0.5% or 1% sodium thiosulfate. These results indicate that the MEM cell growth medium comprising 0.5% or 1% sodium thiosulfate has an activity of 10-fold lower dilution than when only MEM cell culture medium is used.

Determination of the HAdV infectious titers

HAdV infectious titers (TCID₅₀) were determined using a fixed virus copy number of 1×10^5 for inoculation. After approximately 14 days of culture, the cultures were inspected to observe the CPE around the dilution limit (at line 16) to determine TCID₅₀; the results are presented in Table 1. For HAdV-54, the CPE observed on day 17 of culture was used for calculation because this type showed a significant growth delay. Although there was some variability among the strains, the infectious titers were within a range of 4.06–4.67 log₁₀ TCID₅₀/ml (Table 1).

Antiseptic activity of ALTANT for several HAdV types

The results did not differ considerably between the two independent assays. The

antiseptic activities were determined based on EN-14476 of CEN, and the results are summarized in Table. On day 14, after HAdV inoculation, the complete virucidal effect was observed for all adenovirus types when they were allowed to react with ALTANT for 3 minutes. Even in 1 minute, ALTANT satisfied the EN-14476 criterion for the antiviral effect ($> 4\log_{10}$) for all HAdV types tested in the present study. For PCF-related HAdV types, i.e., HAdV-1, HAdV-2, HAdV-3, HAdV-4, HAdV-5, and HAdV-6), ALTANT showed 80.5%–99.22% virucidal effects even in a short reaction time (flash; within 3 seconds from ALTANT-HAdV reaction to neutralization). HAdV-7 associated with severe respiratory disease showed high susceptibility to the ALTANT treatment. HAdV-11 associated with HC also showed a high susceptibility even in a short reaction time. Among EKC-associated HAdV types, HAdV-37, HAdV-53, HAdV-54, HAdV-56, HAdV-64, and HAdV-85 were less susceptible than PCF in flash, with only 50% and 75% virucidal effects being observed for HAdV-53 and HAdV-64, respectively, even after 10 seconds of reaction. However, ALTANT showed approximately 90% virucidal effects on HAdV-37, HAdV-54, HAdV-56, and HAdV-85, which increased to $\geq 99\%$ in 30 seconds for all the types tested.

Discussion

The drug susceptibilities for different types of HAdV varies according to the type of HAdV (22,24) and should be tested using clinically important types of HAdV. HAdV is highly resistant to common hand sanitizers such as ethanol (2,3,22–24). Ozone is a strong oxidizer and exhibits a very high potency and disinfecting efficacy. While the stable use of ozone has been difficult, ALTANT is a formulation wherein ozone is stabilized in an alcoholic solution for more than 3 years (<http://www.e-teck.co.jp/>). ALTANT is cytotoxic to cultured cells, but we demonstrated that its cytotoxicity could be neutralized by 0.5% sodium thiosulfate. In the evaluation of disinfectants against bacteria according to the standardized reference EN1276 of CEN, common neutralizers are shown; however, in the case of viruses, many of these culture medium components affect the growth of host cells and could not be used (data not shown). Therefore, sodium thiosulfate alone was used to test the neutralizing effect, as described in the previous reports (24).

After the drug was allowed to react with the virus for varying lengths of time, the reaction mixtures were subjected to 2-fold serial dilutions, and the neutralizing effect was determined by measuring the virus decrease rates using the CPE occurrence as an indicator after culturing for 7-21 days. The assay used in this study allowed us to determine the endpoint of the complete HAdV-disinfecting effect by comparing the result

with that of the control group. Generally, TCID₅₀, PFU, and virus copy numbers are used to indicate infectious titers of viruses. When a single type of virus is used in the assay, one method is usually enough to determine the infectious titer depending on a condition. When multiple types of viruses are used in the assay, it is difficult to set a common infectious titer because their growth conditions and capacities vary. Different HAdV types grow at varying rates (30) and differ in the drug susceptibility (2,23,24), and thus it is difficult to determine infectious titers of such viruses. Accordingly, the observation was performed up to day 21, and the infectious titer of each HAdV type was determined during this period. For the evaluation of ALTANT, the percentage of viral CPE decrease for each HAdV type was calculated in reference to the control group (no drug reaction). Moreover, as the reacted virus solution in this method is subjected to limiting dilution, and CPE caused by the diluted virus is observed, the virus decrease rate at the highest dilution indicates the virucidal effect. The sterilizing antiviral effect was determined using the longest possible length of the observation period. In CPE assays for adenoviruses in 96-well plates, the CPE in one well equals 1 CFU, particularly for HAdV-5. Mathematically, lanes 16–17 correspond to 1 virus copy. For virtually all strains, lanes 14–16 were limits of CPE detection. In this study, the virus nucleic acid copy number and infectious titer were approximately in agreement for some strains or differed by about 10-fold for some

other strains. This finding is consistent with previous reports (30). The isopropanol solution, which is the base of ALTANT, showed no anti-adenovirus effect (data not shown). ALTANT showed efficacy levels below $4\log_{10}$, which is regarded as the efficiency limit by EN-14476. 70% ethanol or isopropanol with/without 0.5% chlorhexidine-digluconate showed no disinfection efficacy for HAdV with reduction levels below $4\log_{10}$ (24).

In general, disinfectants are known to show large activity changes, decreases in most cases, in actual settings where viruses or bacteria of interest present, because of influences from many coexisting substances, such as proteins and salts (22). In this study, the drug effect was evaluated in the presence of many serum components which reduce the effectiveness of disinfectants (a condition closer to the actual environment). However, the effects in actual foods and environments were not evaluated and remain to be tested in the future. ALTANT is not approval usage to the mucous membrane, however, there is no corrosion or discoloration of the surface of human skin or environmental substances as with alcohol sterilization preparation (<http://www.e-teck.co.jp/>). ATLANT is used by direct-spray or indirectly using a soft cloth such as cotton gauze for the surface of hands and environmental disinfection(http://www.e-teck.co.jp/?page_id=171)

In this study, among PCF-related HAdV types, HAdV-3 was found to be slightly less

susceptible to ALTANT. EKC-associated strains showed characteristic responses for low sensitivity within 3 - 10 seconds, presumably because of a unique mechanism that they use to enter host cells (5,31). Since all HAdV types tested were highly susceptible to ALTANT, the sufficient anti-HAdV effect can be expected in simple hand sanitization and environmental disinfection using ALTANT.

Acknowledgement

This study was partly supported by a Ministry of Health, Labour and Welfare Grant in Aid for Scientific Research (16809509).

Conflict of interest

This study was conducted as part of a collaborative research agreement between the National Institute of Infectious Diseases and E-TECH Corporation. The author; Eiichi Yoshida is CEO of E-TECH Corporation. All authors declare no conflicts of interest.

References.

1. Gonzalez G, Yawata N, Aoki K, et al. Challenges in management of epidemic

- keratoconjunctivitis with emerging recombinant human adenoviruses. *J Clin Virol.* 2019;112:1–9.
2. Sauerbrei A, Sehr K, Brandstädt A, et al. Sensitivity of human adenoviruses to different groups of chemical biocides. *J Hosp Infect.* 2004;57(1):59–66.
 3. Wutzler P, Sauerbrei A. Virucidal efficacy of a combination of 0.2% peracetic acid and 80% (v/v) ethanol (PAA-ethanol) as a potential hand disinfectant. *J Hosp Infect.* 2000;46(4):304–8.
 4. Takahashi K. Useful manifestations to detect adenovirus in children with upper respiratory infections : A retrospective study. 2019;(January):1–5.
 5. Hashizume M, Aoki K, Ohno S, et al. Disinfectant potential in inactivation of epidemic keratoconjunctivitis-related adenoviruses by potassium peroxymonosulfate. *Eur J Ophthalmol.* 2019 Dec 9;112067211989140.
 6. Mori Y, Miyamoto T, Kato K, et al. Different Risk Factors Related to Adenovirus- or BK Virus-Associated Hemorrhagic Cystitis following Allogeneic Stem Cell Transplantation. *Biol Blood Marrow Transplant.* 2012;18(3):458–65.
 7. Umekawa T, Kurita T. Acute Hemorrhagic Cystitis by Adenovirus Type 11 with and without Type 37 after Kidney Transplantation. *Urol Int.* 1996;56(2):114–6.
 8. Yagisawa T, Nakada T, Takahashi K, et al. Acute Hemorrhagic Cystitis Caused

- by Adenovirus after Kidney Transplantation. *Urol Int*. 1995;54(3):142–6.
9. Shindo K, Kitayama T, Ura T, et al. Acute Hemorrhagic Cystitis Caused by Adenovirus Type 11 after Renal Transplantation. *Urol Int*. 1986;41(2):152–5.
 10. Dailey Garnes NJM, Ragoonanan D, Aboulhosn A. Adenovirus infection and disease in recipients of hematopoietic cell transplantation. Vol. 32, *Current Opinion in Infectious Diseases*. 2019. p. 591–600.
 11. Lee YJ, Chung D, Xiao K, et al. Adenovirus viremia and disease: comparison of T cell-depleted and conventional hematopoietic stem cell transplantation recipients from a single institution. *Biol Blood Marrow Transplant*. 2013 Mar ;19(3):387–92.
 12. Sahin U, Toprak SK, Atilla PA, et al.. An overview of infectious complications after allogeneic hematopoietic stem cell transplantation. Vol. 22, *Journal of Infection and Chemotherapy*. 2016. p.505–14.
 13. Legoff J, Feghoul L, Mercier-Delarue S, et al. Broad-range PCR-electrospray ionization mass spectrometry for detection and typing of adenovirus and other opportunistic viruses in stem cell transplant patients. *J Clin Microbiol*. 2013 Dec 1;51(12):4186–92.
 14. Sedláček P, Petterson T, Robin M, et al. Incidence of Adenovirus Infection in

- Hematopoietic Stem Cell Transplantation Recipients: Findings from the AdVance Study. *Biol Blood Marrow Transplant*. 2019 Apr;25(4):810–8.
15. Hubmann M, Fritsch S, Zoellner AK, et al. Occurrence, risk factors and outcome of adenovirus infection in adult recipients of allogeneic hematopoietic stem cell transplantation. *J Clin Virol*. 2016;82:33–40.
 16. Pochon C, Voigt S. Respiratory virus infections in hematopoietic cell transplant recipients. *Front Microbiol*. 2019;10 (Jan):1–17.
 17. Hanaoka N, Ito S, Konagaya M, Nojiri N, et al. Infectious human adenoviruses are shed in urine even after disappearance of urethral symptoms. *PLoS One*. 2019;14(3):1–18.
 18. Enomoto M, Fujimoto T, Konagaya M, et al. Cultivation for 21 days should be considered to isolate respiratory adenoviruses from samples containing small numbers of adenoviral genomes. *Jpn J Infect Dis*. 2010 Sep;63(5):338–41.
 19. Seidler V, Linetskiy I, Hubálková H, et al. Ozone and its usage in general medicine and dentistry. A review article. *Prague Med Rep* 2008;109(1):5–13.
 20. Kosaka K, Yamada H, Shishida K, et al. Evaluation of the treatment performance of a multistage ozone/hydrogen peroxide process by decomposition by-products. *Water Res*. 2001 Oct;35(15):3587–94.

21. Thurston-Enriquez JA, Haas CN, Jacangelo J, et al. Inactivation of enteric adenovirus and feline calicivirus by ozone. *Water Res.* 2005 Sep;39(15):3650–6.
22. Kampf G. Efficacy of ethanol against viruses in hand disinfection. *J Hosp Infect.* 2018 Apr;98(4):331–8.
23. Iwasawa A, Niwano Y, Kohno M, et al. Virucidal Activity of Alcohol-Based Hand Rub Disinfectants. *Biocontrol Sci.* 2012;17(1):45–9.
24. Uzuner H, Karadenizli A, Er DK, et al. Investigation of the efficacy of alcohol-based solutions on adenovirus serotypes 8, 19 and 37, common causes of epidemic keratoconjunctivitis, after an adenovirus outbreak in hospital. *J Hosp Infect.* 2018;100(3):e30–6.
25. Fujimoto T, Hanaoka N, Konagaya M, et al. Evaluation of a silver - amplified immunochromatography kit for adenoviral conjunctivitis. *J Med Virol.* 2019; 91: 1030 - 1035.
26. Watanabe M, Kohdera U, Kino M, et al. Detection of adenovirus DNA in clinical samples by SYBR Green real-time polymerase chain reaction assay. *Pediatr Int.* 2005 Jun;47(3):286–91.
27. Kärber G. Beitrag zur kollektiven Behandlung pharmakologischer Reihenversuche. *Naunyn Schmiedebergs Arch Exp Pathol Pharmacol.* 1931

Jul;162(4):480–3.

28. Spearman CE. Review of The Method of “Right and Wrong Cases” (‘Constant Stimuli’) without Gauss’s Formula. *Psychol Bull.* 1909 Jan;6(1):27–8.
29. Romanowski E.G., Yates K.A., Connor K.E.O., et al. The Evaluation of Polyhexamethylene Biguanide (PHMB) as a Disinfectant for Adenovirus. *JAMA Ophthalmol.* 2013;131:495–498.
30. Kaneko H, Suzutani T, Aoki K, et al. Epidemiological and virological features of epidemic keratoconjunctivitis due to new human adenovirus type 54 in Japan. *Br J Ophthalmol.* 2011;95(1):32–6.
31. Stasiak AC, Stehle T. Human adenovirus binding to host cell receptors: a structural view. *Med Microbiol Immunol.* 2019 Nov 29;(0123456789).

Table. Adenovirus reduction efficacy of ALTANT*

HAdV type	HAdV titer# Log ₁₀ TCID ₅₀ / ml	< 3"		10"		30"		1'		3'		5'	
		Log ₁₀	%	Log ₁₀	%	Log ₁₀	%	Log ₁₀	%	Log ₁₀	%	Log ₁₀	%
1	4.67	1.20	93.75	1.8	98.44	2.71	99.80	3.61	99.98	<4.67	<99.99	<4.67	<99.99
2	4.67	1.81	98.44	1.8	98.44	2.71	99.80	3.31	99.95	<4.67	<99.99	<4.67	<99.99
3	4.06	0.90	87.50	1.2	93.75	2.11	99.22	3.61	99.98	<4.06	<99.99	<4.06	<99.99
4	4.06	2.11	99.22	2.4	99.61	2.71	99.80	<4.06	<99.99	<4.06	<99.99	<4.06	<99.99
5	4.36	1.20	93.75	1.5	96.88	1.81	98.44	3.61	99.98	<4.36	<99.99	<4.36	<99.99
6	4.67	0.90	87.50	1.8	98.44	2.71	99.80	4.21	99.99	<4.67	<99.99	<4.67	<99.99
7	4.06	1.51	96.88	2.4	99.61	3.01	99.90	<4.06	<99.99	<4.06	<99.99	<4.06	<99.99
11	4.36	2.11	99.22	2.4	99.61	2.71	99.80	3.61	99.98	<4.36	<99.99	<4.36	<99.99
37	4.67	0.90	87.50	1.2	93.75	3.61	99.98	<4.67	<99.99	<4.67	<99.99	<4.67	<99.99
53	4.67	0.30	50.00	0.3	50.00	3.01	99.90	<4.67	<99.99	<4.67	<99.99	<4.67	<99.99
54\$	4.06	0.60	75.00	1.2	93.75	2.71	99.80	<4.06	<99.99	<4.06	<99.99	<4.06	<99.99
56	4.97	0.90	87.50	1.2	93.75	3.31	99.95	<4.97	<99.99	<4.97	<99.99	<4.97	<99.99
64	4.67	0.30	50.00	0.6	75.00	4.21	99.99	<4.67	<99.99	<4.67	<99.99	<4.67	<99.99
85	4.67	0.90	87.50	0.9	87.50	2.41	99.61	<4.67	<99.99	<4.67	<99.99	<4.67	<99.99

*Reaction times are indicated as follows: 10"" means 10 seconds, and 1' means 1minute.

#1 × 10⁵ copy of HAdV was inoculated and incubated until determined titer at day 14.

Virus titer was calculated with CPE observation on day 17 because of the slow growth phenotype of HAdV54.